

Athlete's Heart



- Outline
 - Exercise-induced Cardiac Remodeling
 - Health vs. Disease
 - LV chamber enlargement
 - RV chamber enlargement
 - LV wall thickening

Historical Overview

1899: Initial observations by Henschen and Darling

- Cardiac enlargement by physical exam

100+ years of scientific study:





AMERICAN
COLLEGE *of*
CARDIOLOGY

Cardiac Remodeling

Hemodynamic Stress of Sport



Pathophysiology of Disease



Cardiac Remodeling: *Exercise*

Endurance Activities (Isotonic)



Sustained \uparrow CO

- 4 to 5 times rest
- $\uparrow \uparrow \uparrow$ HR & \uparrow SV
- Vasodilation

Volume Challenge

Strength Activities (Isometric)



Repetitive \uparrow SBP


- Systolic BP $>$ 200 mmHg
- Skeletal Muscle Contraction
- Vasoconstriction

Pressure Challenge



AMERICAN
COLLEGE *of*
CARDIOLOGY


FIGURE Classification of Sports



Levine et al
JACC 2015


AMERICAN
COLLEGE of
CARDIOLOGY

**Normal "Pre-training"
Cardiac Structure and Function**




Endurance
Training

Strength
Training



Characteristic Features

- Eccentric LVH and RV dilation
- Batrial enlargement
- Normal to slightly reduced resting LVEF
- Normal or enhanced early diastolic function



Characteristic Features

- Concentric LVH
- Enlarged left atrium
- Hyperdynamic resting LVEF
- Reduced early diastolic function
- Altered LV mechanics

Weiner & Baggish. *Prog Cardiovasc Dis* 2012;54:380.



**AMERICAN
COLLEGE of
CARDIOLOGY**

Determinants of Myocardial Adaptation

- Sporting discipline
- Gender
 - Females < males
- Ethnicity
 - ↑ wall thickness in Afro-Caribbean descent
- Genetics / Molecular pathways
- Exercise exposure duration and “dose”




AMERICAN
COLLEGE of
CARDIOLOGY


Health vs. Disease

- Can we separate athletic remodeling from pathology?
 - Yes, the majority of the time when a systematic approach is used.

Evaluation “Tool Kit”



LV Chamber Enlargement



Pelliccia et al. *Ann Intern Med* 1999;130:23.



AMERICAN
COLLEGE of
CARDIOLOGY

LV Chamber Enlargement

Table 4 Echocardiographic findings from the study population of university athletes

Parameter	Male (n = 300)		Female (n = 197)	
	Normal (n = 209)	Physiologic remodeling (n = 91)	Normal (n = 178)	Physiologic remodeling (n = 19)
Structural parameters				
Interventricular septal thickness (mm)	9.8 ± 0.9	11.6 ± 0.5	8.3 ± 0.7*	10.6 ± 0.5 [†]
LV posterior wall thickness (mm)	10.0 ± 1.2	11.8 ± 1.4	8.6 ± 1.1*	10.7 ± 0.7 [†]
LV inner dimension at end-diastole (mm)	51 ± 3	57 ± 5	42 ± 4*	54 ± 4 [†]
LA diameter (mm)	36 ± 4	40 ± 4	32 ± 3*	38 ± 4
RV end-diastolic diameter (mm)	30 ± 5	36 ± 3	28 ± 4*	33 ± 3 [†]
Functional parameters				
LV ejection fraction (%)	65 ± 7	58 ± 4	68 ± 6	64 ± 6 [†]
Transmitral E wave (cm/sec)	86 ± 16	96 ± 13	81 ± 17	88 ± 12
Transthoracic area (cm ²)	10.0 ± 1.2	11.8 ± 1.4	8.6 ± 1.1*	10.7 ± 0.7 [†]
E' (cm/sec)	10.0 ± 1.2	11.8 ± 1.4	8.6 ± 1.1*	10.7 ± 0.7 [†]
E' (cm/sec)	10.0 ± 1.2	11.8 ± 1.4	8.6 ± 1.1*	10.7 ± 0.7 [†]
A' (cm/sec)	10.0 ± 1.2	11.8 ± 1.4	8.6 ± 1.1*	10.7 ± 0.7 [†]
A' (cm/sec)	10.0 ± 1.2	11.8 ± 1.4	8.6 ± 1.1*	10.7 ± 0.7 [†]

**25% of US college athletes
exceed gender recommended LVIDd limit**

*P < .05 for comparison with male athletes in the normal cardiac structure and function group.

[†]P < .05 for comparison with male athletes in the physiologic remodeling group.

Weiner et al. *J Am Soc Echocardiogr* 2012;25:568.



AMERICAN
COLLEGE of
CARDIOLOGY

- *Physiologic LV Chamber Enlargement:*

- Expected with endurance training.
- Accompanied by proportionate increase in wall thickening (Eccentric LVH).
- Accompanied by normal to low normal resting LVEF (~50%).
- TDI / Strain assessment with preserved or enhanced function.
- Accompanied by “other” chamber enlargement (RV, LA).
- LVIDd “cut-offs” are not helpful.
- When in doubt, exercise testing is very useful (confirm LV augmentation and document supranormal exercise capacity).



AMERICAN
COLLEGE *of*
CARDIOLOGY

RV Chamber Enlargement




Figure 2 Range of values for RV inflow dimension in endurance athletes ($n = 102$).




Figure 3 Range of values for RV proximal outflow dimension in endurance athletes ($n = 102$).

Oxborough et al. *J Am Soc Echocardiogr* 2012;25:263.



AMERICAN
COLLEGE of
CARDIOLOGY

- *Physiologic RV Chamber Enlargement:*


- Expected with endurance training.
- Global RV process without sacculation, aneurysmal dilation, segmental dysfunction, or fibrosis (?).
- RV dimensions “cut-offs” are not helpful.
- “Always” associated with LV remodeling (concomitant LV enlargement but no RVH).
- Accompanied by normal to low normal resting FAC / RVEF.
- TDI / Strain assessment with preserved or enhanced function.
- If in doubt, comprehensive exercise testing and rhythm monitoring.



AMERICAN
COLLEGE *of*
CARDIOLOGY


Thick LV Walls

Adult Athletes



Pelliccia et al. *N Engl J Med* 1991.

Junior Athletes



Sharma et al. *J Am Coll Cardiol* 2002.

Least frequent, but most problematic



AMERICAN
COLLEGE of
CARDIOLOGY

Thick LV Walls



Table 4 Echocardiographic findings from the study population of university athletes

Parameter	Male (n = 300)		Female (n = 197)	
	Normal (n = 209)	Physiologic remodeling (n = 91)	Normal (n = 178)	Physiologic remodeling (n = 19)
Structural parameters				
Interventricular septal thickness (mm)	9.8 ± 0.9	11.6 ± 0.5	8.3 ± 0.7*	10.6 ± 0.5 [†]
LV posterior wall thickness (mm)	10.0 ± 1.2	11.8 ± 1.4	8.6 ± 1.1*	10.7 ± 0.7 [†]
LV inner dimension at end-diastole (mm)	51 ± 3	57 ± 5	42 ± 4	54 ± 4 [†]
LA diameter (mm)	36 ± 4	40 ± 4	32 ± 3*	38 ± 4
RV end-diastolic diameter (mm)	30 ± 5	36 ± 3	28 ± 4*	33 ± 3 [†]
Functional parameters				
LV ejection fraction (%)	65 ± 7	58 ± 4	68 ± 6	64 ± 6 [†]
Transmitral E wave (cm/sec)	86 ± 16	96 ± 13	81 ± 17	88 ± 12
Transthoracic aortic diameter (cm)	3.1 ± 0.2	3.4 ± 0.2	2.8 ± 0.2	3.2 ± 0.2 [†]
LA, LV, RV, and RA volumes (L)				
LA	100 ± 10	110 ± 10	80 ± 10	100 ± 10 [†]
LV	150 ± 10	160 ± 10	120 ± 10	140 ± 10 [†]
RV	100 ± 10	110 ± 10	80 ± 10	100 ± 10 [†]
RA	100 ± 10	110 ± 10	80 ± 10	100 ± 10 [†]

**Not a single healthy college athlete
with walls > 14 mm**

*P < .05 for comparison with male athletes in the normal cardiac structure and function group.

[†]P < .05 for comparison with male athletes in the physiologic remodeling group.



AMERICAN
COLLEGE of
CARDIOLOGY

Weiner et al. *J Am Soc Echocardiogr* 2012;25:568.

- *Physiologic Thick LV Walls:*

- Physiologic concentric LVH is symmetric without regional variation.
 - Marked asymmetry is pathology until proven otherwise.
- Wall thickness “cut-offs” are VERY helpful.
- Accurate absolute thicknesses >15 mm are pathologic until proven otherwise.
- E' values may be helpful, but not diagnostic
- Exercise testing (CPET) is a useful discriminator
- Detraining may be necessary to arrive at a final diagnosis.

This is the HCM mimicker



AMERICAN
COLLEGE of
CARDIOLOGY

Other Areas of Study

- Atria
 - LA dilation: endurance > strength athletes
- LA function: atrial strain and contraction
- Aorta
 - Sinus of Valsalva: 3.2 mm greater in athletes

Iskander et al. *JACC Cardiovasc Imaging*, 2015.


Iskander et al. *Circulation*, 2013.



AMERICAN
COLLEGE of
CARDIOLOGY

Future Directions

- Myocardial mechanics
 - LV strain, twist (regional function)
- Cardiac MRI



La Gerche et al. *Eur Heart J.* 2012; 33:998.

Summary



Thick LV Walls



Key Differential Diagnosis

Hypertrophic cardiomyopathy
Hypertensive heart disease
Infiltrative heart disease
Valvular heart disease

Clinical Factors c/w of Athlete's Heart

Strength training background
No subjective symptoms
Benign family history
Normal subjective exercise capacity

Echo Findings c/w Athlete's Heart

Mild symmetric LVH (walls <15 mm)
Normal RV dimensions
Normal / mildly enlarged LA
Normal aortic valve function
Normal mitral valve anatomy

Additional Diagnostic Considerations

Exercise testing (VO₂ assessment)
24h ambulatory BP monitor
Cardiac MRI
? Prescribed detraining

Dilated LV Chamber



Key Differential Diagnosis

Idiopathic dilated cardiomyopathy
Toxic (ETOH, drugs) cardiomyopathy
Infectious cardiomyopathy
Cardiomyopathy 2° tachyarrhythmia

Clinical Factors c/w of Athlete's Heart

Endurance training background
No subjective symptoms
Benign family history
No history of prior illness / substance abuse
Normal subjective exercise capacity

Echo Findings c/w Athlete's Heart

Concomitant RV dilation
Mild LV wall thickening
Supra-normal LV diastolic indices
Normal / mildly enlarged LA & RA

Additional Diagnostic Considerations

Exercise testing (VO₂ assessment)
Ambulatory rhythm monitoring
Cardiac MRI

Dilated RV Chamber



Differential Diagnosis

Arrhythmogenic RV cardiomyopathy
Idiopathic dilated cardiomyopathy
Pulmonary HTN / congenital heart disease
Sarcoidosis
Cardiomyopathy 2° tachyarrhythmia

Clinical Factors c/w of Athlete's Heart

Endurance training background
No subjective symptoms
Benign family history
Normal subjective exercise capacity

Echo Findings c/w Athlete's Heart

Concomitant LV dilation
Normal RV morphology
Supra-normal LV diastolic indices
Normal / mildly enlarged LA & RA
Normal RV systolic pressure

Additional Diagnostic Considerations

Signal averaged ECG
Exercise testing (VO₂ assessment)
Ambulatory rhythm monitoring
Cardiac MRI